

# Possible detection of a magnetic field in T Tauri

D.A. Smirnov<sup>1</sup>, S.N. Fabrika<sup>2</sup>, S.A. Lamzin<sup>1,3</sup>, and G.G. Valyavin<sup>2</sup>

<sup>1</sup> Sternberg Astronomical Institute, Moscow V-234, 119992, Russia

<sup>2</sup> Special Astrophysical Observatory, Nizhnij Arkhyz, 357147, Russia

<sup>3</sup> Isaak Newton Institute of Chile Branch in Moscow, Russia

Received / Accepted

**Abstract.** Medium-resolution ( $R \simeq 15000$ ) circular spectropolarimetry of T Tauri is presented. The star was observed twice: on November 11, 1996 and January 22, 2002. Weak circular polarization has been found in photospheric absorption lines, indicating a mean surface longitudinal magnetic field  $B_{\parallel}$  of  $160 \pm 40$  G and  $140 \pm 50$  G at the epoch of the first and second observations respectively. While these values are near the detection limit of our apparatus, we believe that they are real. In any case one can conclude from our data that  $B_{\parallel}$  of T Tau does not significantly exceed 200 G, which is much less than surface magnetic field strength of the star ( $> 2.3$  kG) found by Guenther et al. (1999) and Johns-Krull et al. (2000). We discuss possible reasons of this difference.

**Key words.** stars: pre-main sequence – stars: magnetic fields – stars: individual: T Tau

## 1. Introduction

Classical T Tauri stars (CTTSs) are low mass pre-main sequence stars which have accretion disks (Bertout, 1989). Magnetic fields are believed to play a crucial role in the evolution of CTTSs angular momentum and in the interaction between the central star and the circumstellar matter. Surface magnetic fields  $\sim 1$  kG are expected from theoretical arguments for these objects – see e.g. Shu et al. (2000) and references therein.

Meanwhile, no field has been directly detected in CTTSs until recently although several attempts have been made (Babcock 1958; Brown & Landstreet 1981; Johnstone & Penston 1986, 1987). Recently two groups independently succeeded in detecting magnetic fields on T Tau, LkCa 15 (Guenther et al., 1999; Johns-Krull et al., 2000) and BP Tau (Johns-Krull et al., 1999a,b).

Two general methods have been used to measure the magnetic field of T Tau. The first one is based on the fact that magnetic fields produce an enhancement of line equivalent widths that vary with magnetic sensitivity (Lande factor and line splitting pattern). With this technique one can derive the product of the surface magnetic field strength,  $B$ , and surface filling factor  $f$ , i.e.  $B \cdot f$ -value – see Basri et al. (1992) for details. In the case of T Tau, for example, Guenther et al. (1999) found  $B \cdot f = 2.35 \pm 0.15$  kG, which means the average value of  $B$  itself is  $\geq 2.3$  kG.

The other direct method for detecting magnetic fields is to look for net circular polarization in Zeeman-sensitive

lines. Generally, Zeeman  $\sigma$  components are elliptically polarized, with the components of opposite helicity split to either side of the nominal wavelength  $\lambda_0$ . If there is a net longitudinal component of the magnetic field on the stellar surface, net polarization results in a shift between Zeeman-sensitive lines observed in right- and left-circularly polarized (RCP and LCP) light. The shift is

$$\Delta\lambda_{rl} \simeq 9.3 \cdot 10^{-7} g \lambda_0^2 B_{\parallel}, \quad \text{mÅ} \quad (1)$$

where  $g$  is the Lande factor of the transition, which we adopt from the VALD database (Kupka et al. 1999). Here the mean longitudinal magnetic field averaged over the star including limb darkening,  $B_{\parallel}$ , and  $\lambda_0$  are expressed in kG and Å respectively.

Only upper limits of  $B_{\parallel}$  were found in the case of T Tau with this technique: 1000 G (Babcock, 1958) and 810 G (Brown & Landstreet, 1981) at the  $3\sigma$  level. We present here results of our more sensitive spectropolarimetric measurements of T Tau's longitudinal magnetic field. The following set of T Tau's atmospheric parameters were adopted:  $T_{eff} \simeq 5250$  K, (K0 V spectral type),  $\log g = 3.73$  (White & Ghez, 2001) and solar metallicity.

## 2. Observations

We observed T Tau with the 6 m telescope of the Special Astrophysical Observatory on November 21, 1996 and January 22, 2002. The Main Stellar Spectrograph (MSS - Panchuk, 2001) equipped with the polarimetric analyzer, (Najdenov & Chountonov, 1976; Chountonov, 1997) was

used. The analyzer splits the incoming beam into two parallel beams that are offset from one another along the spectrograph slit by 5 arcseconds. One beam contains the RCP component of the spectrum and the other the LCP component. As a result, two spectra with opposite circular polarization  $\sim 200 \text{ \AA}$  in length were projected onto an  $1160 \times 1024$  pixel CCD. The spectral band  $6300 - 6500 \text{ \AA}$  was observed in 1996 and  $6600 - 6800 \text{ \AA}$  in 2002. The width of MSS's slit was  $0.5''$  providing a 2-pixel spectral resolution of  $R = \lambda/\Delta\lambda \simeq 15000$ .

Spectra were processed as follows. Dark current, night sky emission and detector bias as well as cosmic ray traces, were removed in a standard way, using routines from the MIDAS software package. A spectrum of a thorium-argon lamp was used for wavelength calibration. Then we identified spectral lines adopting line information from the VALD database (Kupka et al., 1999). The shifts between positions of identical lines in spectra with opposite circular polarization in the same image,  $\Delta\lambda_{rl}$ , were derived by means of a Fourier cross-correlation method. The method appears to be relatively insensitive to uncertainties in the continuum level position and gives the ability to evaluate the displacement of line profiles as a whole (Klochko et al., 1996).

We took into account systematic instrumental effects (SIEs) in different ways reducing the 1996 and 2002 observations. In the first case, we used observations of the G9.5 III star  $\varepsilon$  Tau, which presumably has negligible field. Specifically, we assumed that nonzero values of  $\Delta\lambda_{rl}(\lambda)$  in  $\varepsilon$  Tau's spectrum result from instrumental effects only and used them to correct  $\Delta\lambda_{rl}$ -values in the spectrum of T Tau – see Romanyuk et al. (1998) for more details. For this reason, our November 21, 1996 observations were organized in the following way: initially we observed  $\varepsilon$  Tau, then T Tau and then  $\varepsilon$  Tau again. Approximate signal-to-noise ratio (S/N) of individual spectra, number of observed spectra,  $n$ , visual magnitude  $V$ , and total exposure time,  $t_e$  (in seconds), for each target are presented in Table 1.

**Table 1.** Journal of Observations

| Date       | Target            | V   | n | $t_e$ | S/N |
|------------|-------------------|-----|---|-------|-----|
| 22.11.1996 | $\varepsilon$ Tau | 3.5 | 5 | 220   | 300 |
|            | T Tau             | 9.9 | 3 | 3600  | 150 |
| 21.01.2002 | 75 Tau            | 5.0 | 3 | 1320  | 300 |
|            | $\varepsilon$ Tau | 3.5 | 3 | 600   | 300 |
|            | T Tau             | 9.9 | 3 | 7200  | 100 |
| 24.01.2002 | HD 30466          | 7.3 | 1 | 1400  | 200 |

In the case of the 2002 observations, SIEs were removed as follows. Between exposures the phase compensator was advanced  $1/2$  wave in order to reverse the sense of circular polarization recorded in the two spectra. For each target, the wavelength of a given line in the RCP and

LCP spectra is recorded in the first exposure ( $\lambda_1^r, \lambda_1^l$ ) and in the second ( $\lambda_2^r, \lambda_2^l$ ). Then we calculated

$$\Delta\lambda_{rl} \equiv \lambda^r - \lambda^l = \frac{(\lambda_1^r - \lambda_1^l) + (\lambda_2^r - \lambda_2^l)}{2},$$

which is expected to be free from the main instrumental effects – see Johns-Krull et al. (1999b) for details.


Information on the January 2002 observations is also presented in Table 1, where now  $n$  stands for the number of pairs of individual observations of any particular star, between which the phase compensator was rotated.

### 3. Results

#### 3.1. 1996 observations

Figure 1 presents values of  $\Delta\lambda_{rl}(\lambda)$  derived from all the  $\varepsilon$  Tau spectra observed immediately before and after observations of T Tau. The  $\Delta\lambda_{rl}(\lambda)$  dependence was approximated with a linear function  $\Delta\lambda_{rl} = a \cdot N + b$ , where  $N$  is the pixel number, for both observations. We found  $a = (-2.8 \pm 2.6) \cdot 10^{-3} \text{ m\AA}$  per pixel,  $b = 10.5 \pm 1.4 \text{ m\AA}$  for the open squares in Figure 1 and  $a = (-5.3 \pm 3.4) \cdot 10^{-3} \text{ m\AA}$  per pixel,  $b = 14.4 \pm 1.9 \text{ m\AA}$  for black triangles in Figure 1.

The difference between the parameters of the fits for both datasets (squares and triangles) is within  $2.5\sigma$ , i.e. is not significant from a statistical viewpoint. So one can use the resulting linear function, derived from all points, to remove SIEs from  $\Delta\lambda_{rl}$ -values in the spectra of T Tau. We found  $a = (-4.1 \pm 2.2) \cdot 10^{-3} \text{ m\AA}$  per pixel,  $b = 12.4 \pm 1.2 \text{ m\AA}$  in this case – solid line in Fig.1



**Fig. 1.**  $\Delta\lambda_{rl}$ -values measured in  $\varepsilon$  Tau's spectra observed before (filled triangles) and after (open squares) T Tau. See text for details.

But the  $a$ -value differs from zero by less than  $2\sigma$ , which means that it is possible that the SIEs do not vary along the CCD. With this hypothesis we found the mean shifts are  $11.8 \pm 0.9$ , and  $9.1 \pm 0.7 \text{ m\AA}$  for filled triangles and open squares respectively. The mean shift  $C$  derived for all points is  $10.4 \text{ m\AA}$  – dashed line in Fig.1 – and standard error is  $\sigma_C = 0.8 \text{ m\AA}$ , or  $\sigma_C = 21 \text{ G}$ , assuming  $g = 1$  and  $\lambda_0 = 6400 \text{ \AA}$  according to equation (1). The mean shifts

for filled triangles and open squares differ from  $C$  less than  $2\sigma_C$ . So it appears resonable to use the mean  $C$  to remove SIEs, subtracting it from the measured wavelength shifts between right and left circularly polarized spectra of T Tau.

The corrected  $\Delta\lambda_{rl}$ -values in T Tau's spectra were used to calculate  $B_{\parallel}$  from equation (1). Resulting values of  $B_{\parallel}$  are presented in Table 2 for each of three observed spectra – the last three columns of the table. In some cases, a given line profile was distorted due to cosmic ray events and thus was not used; these cases are marked with a dash. In fact, almost all absorption features in T Tau's spectra are blends of some number of lines, given in the first two columns. So we use an effective Lande factor,  $g_{eff}$ :

$$g_{eff} = \frac{\sum d_i g_i}{\sum d_i},$$

derived from averaging over individual lines which form the blend. Here  $g_i$  is the Lande factor of each line and  $d_i$  is its central depth. This approach appears resonable as long as the Zeeman splitting is less than the spectral resolution of our instrument. We adopted T Tau's  $d_i$ -values from the VALD database specifying  $\log g$  and  $T_{eff}$  of the star as input parameters. Averaging over all lines gives  $B_{\parallel} = +162$  G with standard error  $\sigma_B = 30$  G due to measurment uncertainty. Adding the error due to the uncertainty in  $C$ , the total uncertainty is:

$$\sigma = \sqrt{\sigma_C^2 + \sigma_B^2} \simeq 37 \text{ G}.$$

Thus, we estimate  $B_{\parallel} = +160 \pm 40$  G from our 1996 observations.

### 3.2. 2002 observations

The  $\Delta\lambda_{rl}(\lambda)$ -values for all sets of  $\varepsilon$  Tau's spectra are shown in Figure 2. It can be seen that there is no statistically significant trend in these data around an average value  $\Delta\lambda_{rl} = -0.02 \pm 0.32$  mÅ. Assuming  $g_{eff} = 1$  and  $\lambda_0 = 6700$  Å (the average wavelength of the spectral band), this value corresponds to  $B_{\parallel} = -0.5 \pm 8$  G. Small values ( $\Delta\lambda_{rl} = -0.93 \pm 0.43$  mÅ or  $B_{\parallel} = -20 \pm 10$  G) were also found for an additional giant star 75 Tau (K2 III). Thus, we conclude that the adopted method of SIEs removal works well and confirmed *a posteriori* our choice of  $\varepsilon$  Tau as a zero-field standard during the 1996 observations.

To illustrate the reliability of our technique we included the magnetic star HD30466 in the target list of our 2002 observations. Babcock (1958) found  $B_{\parallel} = +2320 \pm 340$  G and  $B_{\parallel} = +1890 \pm 130$  G for HD30466 from two independent observations. Profiles of the Cr I 6661.08 line in our HD30466's RCP and LCP spectra are shown in Fig.3: they are clearly shifted relative each other ( $\Delta\lambda_{rl} \simeq 126$  mÅ), indicating  $B_{\parallel} \simeq +2.0$  kG, adopting  $g = 1.48$  (Kupka et al. 1999)

As can be seen from Table 3, relative shifts of lines between spectra with opposite polarizations in T Tau's


**Table 2.** Results of 21.11.1996 observations of T Tau

| Ion  | $\lambda$ | $g_{eff}$ | $B_1$ | $B_2$ | $B_3$ |
|------|-----------|-----------|-------|-------|-------|
| Ni I | 6314.65   | 1.18      | –     | +203  | +11   |
| Ni I | 6314.66   |           |       |       |       |
| Fe I | 6315.31   |           |       |       |       |
| Fe I | 6315.42   |           |       |       |       |
| Fe I | 6315.81   |           |       |       |       |
| Fe I | 6318.02   | 0.72      | –     | +638  | –159  |
| Ca I | 6318.11   |           |       |       |       |
| Ni I | 6322.16   | 1.39      | –     | +251  | +88   |
| Fe I | 6322.68   |           |       |       |       |
| Ni I | 6327.59   | 1.05      | –14   | +157  | –     |
| Cr I | 6330.09   | 1.64      | +262  | –36   | +177  |
| Fe I | 6330.85   |           |       |       |       |
| Fe I | 6338.88   | 1.23      | +500  | +200  | –123  |
| Ni I | 6339.11   |           |       |       |       |
| Ca I | 6343.31   | 1.09      | +18   | –     | +321  |
| Fe I | 6344.15   |           |       |       |       |
| Fe I | 6355.03   | 0.98      | +470  | +110  | –     |
| Fe I | 6358.63   | 1.49      | +198  | +35   | +242  |
| Fe I | 6358.70   |           |       |       |       |
| Fe I | 6393.60   | 1.15      | +288  | –46   | +128  |
| Fe I | 6392.54   |           |       |       |       |
| Fe I | 6400.00   | 1.35      | +28   | +365  | –26   |
| Fe I | 6400.32   |           |       |       |       |
| Fe I | 6411.11   | 1.49      | +162  | –     | +248  |
| Fe I | 6411.65   |           |       |       |       |
| Fe I | 6412.20   |           |       |       |       |
| Fe I | 6430.85   | 1.24      | –153  | +399  | –215  |
| Ca I | 6439.08   | 1.12      | +210  | +254  | +313  |
| Ca I | 6449.81   | 1.10      | +48   | +8    | +143  |
| Co I | 6450.08   |           |       |       |       |
| Co I | 6450.25   |           |       |       |       |
| Ca I | 6462.57   | 1.17      | +588  | +124  | –158  |
| Fe I | 6462.71   |           |       |       |       |
| Fe I | 6462.73   |           |       |       |       |
| Ca I | 6471.66   | 1.20      | +319  | –53   | +342  |
| Ca I | 6475.24   | 1.63      | +105  | +232  | +430  |
| Fe I | 6475.62   |           |       |       |       |


spectrum are much less, and we find  $B_{\parallel} = +140 \pm 50$  G after averaging over all the data. The uncertainty is calculated by taking the standard error of the individual points.

## 4. Discussion

From observations in 1996 and 2002, using different methods of SIE correction, we find similar ( $+160 \pm 40$  G and  $+140 \pm 50$  G) values of the mean surface longitudinal magnetic field in T Tau. As far as such small field values are close to the detection limit, it is very important to show that the measured wavelength shift is magnetic in origin. To test this we have measured relative shifts of the S II



**Fig. 2.**  $\Delta\lambda_{rl}$ -values derived from three pairs of  $\epsilon$  Tau's spectra observed in 22.01.2002. See text for details.



**Fig. 3.** A part of RCP (thin line) and LCP (solid line) spectra of HD30466

6716.44 Å and 6730.82 Å forbidden emission lines in our 2002 spectra, which presumably originate far from the star in a region with (nearly) zero magnetic field. We found  $\Delta\lambda_{rl} = +0.8 \pm 1.4$  mÅ for these lines which would give  $20 \pm 35$  G for  $g_{eff} = 1$ . This result can be considered as evidence that the measured wavelength shift of the stellar absorption lines is magnetic in origin.

To further test if it is indeed possible to detect such low  $B_{||}$  value from our spectra we performed the following test. The RCP and LCP spectra of 1D stellar atmosphere were calculated with the SYNTHMAG program (Piskunov, 1998), assuming that  $\log g$  and  $T_{eff}$  are equal to that of T Tau and  $B = B_{||} = 200$  G, i.e. with magnetic field lines perpendicular to the stellar surface. Profiles of lines within 6300-6500 Å spectral band were artificially broadened to take into account stellar rotation and spectral resolution and binned to the appropriate dispersion. Each of these artificial spectra were used to produce three new ones by means of adding noise at the appropriate level. Finally we derived three pairs of RCP and LCP spectra with S/N-ratio near 100, as in the case of our 1996 and/or 2002 observations. By means of cross-correlation of RCP and LCP profiles we then found  $B_{||} = 164 \pm 32$  G in agreement with the initial value of 200 G. Therefore, we believe that we indeed have recorded T Tau's surface magnetic field  $B_{||} \sim 150$  G, in spite of the relatively large scatter of the individual measurements – see Table 2& 3.

**Table 3.** Results of 22.02.2002 observations of T Tau

| Ion  | $\lambda$ | $g_{eff}$ | $B_1$ | $B_2$ | $B_3$ |
|------|-----------|-----------|-------|-------|-------|
| Fe I | 6609.1    | 1.59      | −161  | −33   | −76   |
| Fe I | 6609.7    |           |       |       |       |
| V I  | 6624.8    | 0.94      | +768  | −21   | +918  |
| Fe I | 6625.0    |           |       |       |       |
| Fe I | 6633.4    | 1.48      | +576  | +218  | −299  |
| Fe I | 6633.8    |           |       |       |       |
| Fe I | 6634.1    |           |       |       |       |
| Ni I | 6643.6    | 1.31      | —     | −8    | −182  |
| Fe I | 6663.2    | 1.53      | +160  | +58   | +460  |
| Fe I | 6663.4    |           |       |       |       |
| Fe I | 6677.9    | 1.35      | +143  | −207  | −67   |
| Fe I | 6678.0    |           |       |       |       |
| Al I | 6696.0    | 0.97      | −101  | −182  | +545  |
| Fe I | 6696.3    |           |       |       |       |
| Al I | 6698.7    | 1.25      | +198  | +633  | −214  |
| Fe I | 6699.1    |           |       |       |       |
| Li I | 6707.8    | 1.25      | +79   | +91   | +87   |
| Fe I | 6710.3    | 1.69      | −82   | −162  | −150  |
| Ca I | 6717.7    | 1.01      | +7    | +52   | +232  |
| Ti I | 6743.1    | 1.01      | +735  | +633  | +635  |
| Fe I | 6750.2    | 1.50      | −201  | −77   | +736  |
| Fe I | 6752.7    | 2.01      | −282  | −161  | +61   |
| Ni I | 6767.8    | 1.43      | +192  | −111  | +646  |

Regardless, our observations show T Tau's large scale  $B_{||}$ -value did not significantly exceed 200 G.

This value is much less than the average value of T Tau's surface magnetic field ( $B > 2$  kG) found by Guenther et al. (1999) and Johns-Krull et al. (2000). This will result if there are regions with opposite polarity of magnetic field lines at the stellar surface. This could indicate the magnetic field of T Tau is non-dipolar. Note however, that even in the case of a magnetic dipole, the observed  $B_{||} = 0$ , if the dipole's axis is perpendicular to the line of sight. To demonstrate this more clearly we considered a magnetic dipole with equatorial field strength  $B = 1.6$  kG and calculated using the SYNTHMAG program the relative shifts of RCP and LCP profiles of the Fe I 6430.8 line for different inclination angles  $\beta$  and then derived the "observed" longitudinal field. We found  $B_{||} \simeq 900$ , 400 and 200 G for  $\beta = 0^\circ$ ,  $60^\circ$  and  $75^\circ$  respectively. The mean unsigned surface field values for these inclinations are 2.4, 2.2 and 2.1 kG respectively, i.e. close to values found by Guenther et al. (1999) and Johns-Krull et al. (2000).

T Tau is believed to have a low inclination to its rotation axis based on  $v \sin i$  (Hartmann & Stauffer, 1989) and rotation period measurements (Herbst et al., 1985). Therefore, one can expect that the magnetic axis is also close to the line of sight if it is aligned with or close to the rotation axis. Thus the low value of the average longitudinal magnetic field of T Tau we found can be interpreted

in two possible ways: 1) as an indication of the non-dipole character of the stellar magnetic field, or 2) as evidence of a large ( $\gtrsim 30^\circ$ ) angle between the rotation and magnetic axes.

## 5. Conclusion

Weak circular polarization has been found in photospheric absorption lines of T Tau, indicating a mean longitudinal magnetic field  $B_{\parallel} \sim 150$  G. While this value is near the detection limit of our apparatus, we believe that it is real. In any case one can conclude from our data that  $B_{\parallel}$  on T Tau does not significantly exceed 200 G, i.e. much less than surface magnetic field strength of the star ( $> 2.3$  kG) found by Guenther et al. (1999) and Johns-Krull et al. (2000). This could be the result of the non-dipole character of the magnetic field on T Tau or evidence of a large angle between the rotation and magnetic axes. More precise observations are necessary to distinguish between these possibilities.

*Acknowledgements.* We thank our colleagues G.A. Chountonov, N.E. Piskunov, T.A. Ryabchikova, D.N. Monin and A.A. Panferov for assistance with observations and data reduction as well as helpful discussions. It is a pleasure to thank the referee C.M. Johns-Krull for a careful reading of the manuscript and very useful and benevolent comments. Finally we thank the Russian Fund of Fundamental Research for the grants 02-02-16070 (D.S. and S.L.) and 01-02-16808 (S.F.).

## References

- Babcock H.W., 1958, ApJS, 3, 207  
 Basri G., Marcy G.W., Valenti J.A., 1992, ApJ, 390, 622  
 Bertout C., 1989, ARA&A, 27, 351  
 Brown D.N., Landstreet J.D., 1981, ApJ, 246, 899  
 Chuntunov G.A., 1997, Annual Report of Special Astrophysical Observatory, pp.36 & 56  
 Donati J.-f., Semel M., Carter B.D., Rees D.E., Cameron A.C., 1997, MNRAS 291, 658  
 Guenther, E., Lehmann, H., Emerson, J.P., Staude, J. 1999, A&A, 341 768  
 Hartmann L., Stauffer J.R., 1989, AJ 97, 873  
 Herbst W., Booth J.F., Chugainov P.F. et al., 1985, ApJ 310, L71  
 Johns-Krull, C.M., Valenti, J.A., Hatzes A.P., Kanaan A. 1999a, ApJ, 510, L41  
 Johns-Krull, C.M., Valenti, J.A., Koresko, C. 1999b, ApJ, 516, 900  
 Johns-Krull, C.M. et al., 2000, ASP Conf. Ser. v.223, p.521  
 Johnstone R.M., Penston M.V., 1986, MNRAS 219, 927  
 Johnstone R.M., Penston M.V., 1987, MNRAS 227, 797  
 Klochkova V.G., Yermakov S.V., Panchuk V.E., 1996, Bull. Spec. Astrophys. Obs., 41, 58  
 Kupka F., Piskunov N., Ryabchikova T.A., Stempels H.C., Weiss W.W., 1999, A&AS 138, 119  
 Najdenov I.D., Chuntunov G.A., 1976, Communications of Special Astrophysical Observatory, v.16, p.63 (in Russian)  
 Panchuk V.E., 2001, Preprint of Special Astrophysical Observatory, N 154

- Piskunov, N.E., 1998, in Proc. of the 2nd International Workshop on Solar Polarization, Nagendra K., Stenflo J. (eds.), Bangalore, India, 1998, Kluwer ASSL 243, p. 515  
 Romanyuk I.I., Elkin V.G., Kudryavtsev D.O., Landstreet J.D., Wade G.A., 1998, Bull. Spec. Astrophys. Obs. v.45, p.93  
 Shu, F.H., Najita, J.R., Shang, H., Li, Z.-Y. 2000, in Protostars and Planets IV, eds. V. Mannings, A.P. Boss, & S.S. Russell, University of Arizona Press, Tucson, p. 789  
 White R.J., Ghez A.M., 2001, ApJ 556, 265.